

# EMPIRIC TREATMENT OF COMMUNITY-ACQUIRED PNA IN NON-ICU PATIENTS<sup>^</sup>

Community-acquired pneumonia is defined as pneumonia acquired outside of hospitals.

<b>HMS PREFERRED THERAPY</b> for patients with non-life threatening penicillin allergy (without anaphylaxis, angioedema, hives)	<b>ALTERNATIVE BUT HMS NON-PREFERRED</b> Preferred for patients with cephalosporin allergy, allergy to both macrolides and doxycycline/tetracycline, or severe penicillin allergy.
Ampicillin-Sulbactam 3gm IV q6h <b>OR</b> Ceftriaxone 1gm IV q24h  <b>PLUS</b>  Azithromycin 500mg IV/PO x 1 day, then 250mg q24h x 4 days* <b>OR</b> Clarithromycin 500mg PO BID <b>OR</b> Doxycycline 100mg PO BID	Levofloxacin 750 mg PO/IV Once Daily <b>OR</b> Moxifloxacin 400mg PO/IV Once Daily

\*Consider substituting doxycycline for azithromycin in patients with a macrolide allergy or at risk for prolonged QT interval.

<b>ORAL STEP-DOWN THERAPY WHEN NO ETIOLOGIC PATHOGEN IDENTIFIED FOR CAP**</b>	
Amoxicillin (1g PO 3 x daily) Amoxicillin/clavulanate (2g PO 2 x daily) Cefpodoxime (200mg PO 2 x daily) Cefdinir (300mg PO 2 x daily) Cefditoren (400mg PO 2 x daily) Cefuroxime (500mg PO 2 x daily)	<div style="text-align: center;"> <span style="font-size: 2em;">+ / -</span> </div> Azithromycin, Doxycycline, or Clarithromycin (see dosing above)
<i>Alternatives: Levofloxacin or Moxifloxacin in setting of severe PCN allergy</i>	

<sup>^</sup>Excludes patients with a previous culture positive for MRSA or resistant gram-negative organism in the past year OR patients with severe CAP who were hospitalized and received IV antibiotics in the previous 90 days.

\*\*Suggested dosing only. Please individualize based on renal function or other pertinent clinical factors.

Anaerobic coverage is not routinely warranted in non-critically ill patients with aspiration pneumonia.

For more detail about these guidelines, please see the [Treatment of Community-Acquired Pneumonia Guidelines](#) published by HMS.

# THErapy DURATION & ORAL STEP-DOWN THErapy RECOMMENDATIONS FOR PATIENTS WITH CAP

## DEFINITIONS OF UNCOMPLICATED CAP & COMPLICATED CAP

<p><b>UNCOMPLICATED CAP</b></p>	<p>Patients who do not meet any of the criteria below.</p>
<p><b>COMPLICATED CAP</b></p>	<p>Patients with structural lung disease (e.g. bronchiectasis, pulmonary fibrosis, interstitial lung disease); moderate/severe COPD (excluding COPD exacerbation without pneumonia); documented pneumonia with MRSA, MSSA, or Pseudomonas (or other non-fermenter/gram negative pneumonia); or those who are immunosuppressed.</p>

## DURATION OF ANTIMICROBIAL THERAPY (INCLUDES IV & ORAL)

<p><b>UNCOMPLICATED CAP</b></p>	<p><b>5 Days</b> if patient is afebrile for 48 hours and has <i>no more than one</i> sign of clinical instability* by day 5 of treatment</p> <p>Therapy can be continued for patients who are febrile or clinically unstable* on day 5 of treatment</p>
<p><b>COMPLICATED CAP</b></p>	<p><b>7 Days</b> if patient is afebrile for 48 hours and has <i>no more than one</i> sign of clinical instability* by day 7 of treatment (Note: Azithromycin duration should be no more than 5 days)</p> <p>Therapy can be continued for patients who are febrile or clinically unstable* on day 7 of treatment</p>

\**Signs of Clinical Instability:* O2 saturation < 90% or new oxygen requirement, HR > 100 bpm, RR > 24 bpm, SBP <90 mmHg, altered mental status (different than baseline)

