



# D.I.S.Ch.A.R.G.E.

## Antibiotics: FACTS AND SOLUTIONS



# D.I.S.Ch.A.R.G.E. Antibiotics: FACTS AND SOLUTIONS

1


## Most Antibiotic Stewardship Programs do not Monitor Antibiotic Use at Hospital Discharge

- 8 out of 48 HMS hospitals (HMS Spring Survey 2016)

2

## Half of Antibiotic Exposure Related to Hospitalization Occurs After Discharge (HMS Data)

### HALF OF ANTIBIOTIC USE OCCURS AFTER DISCHARGE



- For some antibiotics, this number is even higher:
  - For example, **65.8%** of fluoroquinolone-days occur after discharge

3

## Without Monitoring, Antibiotic Use at Discharge is Often Excessive (HMS Data)

- For pneumonia, **94.3%** of all excess antibiotic duration occurs after discharge
- Up to half of antibiotics prescribed at discharge have a broad spectrum
  - Pneumonia: **56.5%**
    - Fluoroquinolone: **39.5%** of patients
    - Third-generation cephalosporin: **15.8%** of patients
  - UTI: **31.2%**
    - Fluoroquinolone: **25.4%** (17.3% of asymptomatic bacteriuria)
    - Third-generation cephalosporin: **6.8%** (5.7% of asymptomatic bacteriuria)

# D.I.S.Ch.A.R.G.E. Antibiotics: FACTS AND SOLUTIONS

4

## Inpatient Antibiotic Stewardship May Shift Antibiotic Use to Discharge (HMS data)

- In hospitals actively targeting inpatient fluoroquinolone use:
  - **79.6%** of fluoroquinolone-days occur after discharge
  - Twice as many new fluoroquinolone starts occur at discharge
  - There is less inpatient fluoroquinolone use, but similar discharge and overall days of fluoroquinolone therapy per **1000** patients

5

## Why do Discharge Prescriptions Matter?

- Discharge antibiotics can cause patient harm:
  - Including side effects, Clostridium difficile (CDI), and antibiotic resistance, diarrhea, rash, kidney toxicity, etc.
  - As few as 1-3 days of fluoroquinolone therapy doubles risk of CDI<sup>3</sup>
    - So, changing to fluoroquinolone therapy at discharge puts patients at risk for community-onset CDI
- If we could optimize duration at discharge, we would:
  - **Eliminate 94.3%** of all excess antibiotic duration for pneumonia
  - **Reduce** days of antibiotic therapy by **25%** (2398 days/1000 patients)
    - Half of the reduction would be in broad-spectrum antibiotic therapy

## D.I.S.Ch.A.R.G.E!

*How to improve antibiotic prescribing at hospital discharge.*

### Default orders and order sets

- Consider use of default transitions from IV to oral and recommendations within computerized order-entry to improve early transition to appropriate oral therapy (which can then be continued on discharge)
- Make sure default orders and order sets recommend guideline-appropriate antibiotic choice and duration

### Incentivize

- Consider incorporating discharge antibiotic metrics into quality or compensation targets

### Discharge Summary

- Require documentation of total antibiotic duration in discharge summary
  - Consider enforcing this rule by using smart phrases with hard stops for antibiotic duration in the discharge summary
  - E.g.: To treat (disease), Mr(s) X will continue (abx name) for X additional days, for X days total.

### Checklist

- Use an antibiotic checklist at discharge to evaluate and ensure antibiotic appropriateness

### Audit and Feedback

- Audit and provide feedback of discharge prescriptions (e.g., pharmacists or stewardship team, performance review, quality compensation targets)

Review: Incorporate antibiotic appropriateness into discharge review process using different members of the care team

- For example
  - With pharmacists (when reviewing or filling discharge medications)
  - With bedside nurse (when reviewing discharge medications)
  - During multidisciplinary/discharge rounds

### Guidelines

- Make sure your institutional guidelines include oral antibiotic recommendations for discharge for common infections (e.g., pneumonia, urinary tract infection)
  - Prioritize non-fluoroquinolone antibiotics in guidelines
  - Recommend alternatives to fluoroquinolone antibiotics when possible
  - Provide a recommendation for appropriate duration for different disease states (e.g., 5 days for community-acquired pneumonia), making sure that total duration includes effective inpatient therapy

### Educate providers on guidelines and discharge recommendations

- Formal lectures to residents, physicians (e.g., hospitalist, ID, ED), APPs
- Consider using pocket card
- Consider the use of multiple ways to post guidelines (e.g., websites, apps, printed books)

# D.I.S.Ch.A.R.G.E. Antibiotics: FACTS AND SOLUTIONS

Support for HMS is provided by Blue Cross and Blue Shield of Michigan and Blue Care Network as part of the BCBSM Value Partnerships program. Although Blue Cross Blue Shield of Michigan and HMS work collaboratively, the opinions, beliefs and viewpoints expressed by the author do not necessarily reflect the opinions, beliefs and viewpoints of BCBSM or any of its employees.



Nonprofit corporations and independent licensees of  
the Blue Cross and Blue Shield Association