

Sepsis Team Feedback			
Date of Sepsis Care:			
Patient MRN:			
Time Zero*:			
First Severe Sepsis/Shock BPA:			
Code Sepsis/sepsis alert:			
3-Hour Bundle Elements	Yes/No	Time completed	Goal
Initial Lactate			Drawn within 3 hours of time zero
Blood Cultures			Drawn before antibiotics
Broad Spectrum Antibiotics			Started within 3 hours of Time zero
30cc/kg IVF Bolus: Hypotension or Lactate ≥ 4			Started within 3 hours of Time zero. Required for Hypotension and/or Lactate ≥ 4
6-Hour Bundle Elements			
Repeat Lactate if initial > 2			Drawn within 6 hours of Time zero if initial Lactate > 2
2 BPs Documented within 1 hour of IVF Bolus completion			2 BPs Documented w/i 1 hour of IVF stop time to assess for persistent hypotension
Vasopressor for persistent hypotension after IVF bolus			Patient with persistent hypotension after receiving 30cc/kg IVF bolus
Reassessment of fluid status & tissue perfusion			CVP, ScvO ₂ or Sepsis Reassessment Note completed within 6 hours of time zero
Sepsis Bundle All Requirements Met:			
Comments:			
Opportunities for Improvement:			

* Time zero abstraction is based on CMS abstraction tools for the SEP-1 measure and is intended to give guidance and more real time feedback on team performance. The abstraction is based on best knowledge of the CMS measure, the workflow and the guidance of CMS abstractors.